

Test 2

This test is graded out of 38 marks. No books, notes, unauthorised electronic devices are allowed. You must show all your work, the correct answer is worth 1 mark the remaining marks are given for the work. If you need more space for your answer use the back of the page.

Question 1.¹ (5 marks) A genetic test is used to determine if people have a predisposition for thrombosis, which is the formation of a blood clot inside a blood vessel that obstructs the flow of blood through the circulatory system. It is believed that 3% of people actually have this predisposition. The genetic test is 99% accurate if a person actually has the predisposition, meaning that the probability of a positive test result when a person actually has the predisposition is 0.99. The test is 98% accurate if a person does not have the predisposition. What is the probability that a randomly selected person who tests positive for the predisposition by the test actually has the predisposition?

¹OpenIntro Statistics by D.M. Diez, C.D. Barr and M. Çetinkaya-Rundel, OpenIntro LaTeX, code, and PDFs are released under a Creative Commons BY-SA 3.0 license.

Question 2. The *uniform probability density function* of a continuous random variable X is defined as

$$f(x) = \begin{cases} \frac{1}{b-a} & \text{if } x \in [a, b] \\ 0 & \text{if } x \notin [a, b] \end{cases}$$

- a. (2 marks) Prove that the uniform probability density function is a probability density function.
- b. (2 marks) Find μ and σ for the uniform probability density function.
- c. (2 marks) Express $P(X > \mu \mid X < \mu + \sigma)$ in terms of integrals.

Question 3. A small pond has 11 fish of which 4 are tagged.

- a. (2 marks) If a marine biologist catches 4 fish from the small pond, find a probability function for the discrete random variable X whose values x are the number of tagged fish caught by the marine biologist.
- b. (2 marks) Prove that the function found in part a. is a probability function.
- c. (2 marks) What is the probability that the marine biologist will catch fewer tagged fish than expected.

Question 4.² (5 marks) Suppose a university announced that it admitted 2,500 students for the following year's freshman class. However, the university has dorm room spots for only 1,786 freshman students. If there is a 70% chance that an admitted student will decide to accept the offer and attend this university, what is the approximate probability that the university will not have enough dormitory room spots for the freshman class?

²OpenIntro Statistics by D.M. Diez, C.D. Barr and M. Çetinkaya-Rundel, OpenIntro LaTeX, code, and PDFs are released under a Creative Commons BY-SA 3.0 license.

Question 5.³ If $P(A) = 0.3$, $P(B) = 0.7$ and assuming that events A and B arise from independent random processes,

i. (1 mark) Compute $P(A \text{ and } B)$.

ii. (1 mark) Compute $P(A \text{ or } B)$.

iii. (1 mark) Compute $P(A | B)$.

Question 6. (2 marks) Prove: If A_1, A_2, A_3 are events then $P(A_1 \cap A_2 \cap A_3) = P(A_1)P(A_2 | A_1)P(A_3 | A_1 \cap A_2)$.

Question 7.⁴ Below are final exam scores of 20 Introductory Statistics students.

57, 66, 69, 71, 72, 73, 74, 77, 78, 78, 79, 79, 81, 81, 82, 83, 83, 88, 89, 94

a. (3 marks) The mean score is 77.7 with a standard deviation of 8.44. Use this information to determine if the scores approximately follow the 68 – 95 – 99.7% Rule.

b. (2 marks) What does the Chebyshev Theorem tell us about the data and 2 standard deviation.

³OpenIntro Statistics by D.M. Diez, C.D. Barr and M. Çetinkaya-Rundel, OpenIntro LaTeX, code, and PDFs are released under a Creative Commons BY-SA 3.0 license.

⁴OpenIntro Statistics by D.M. Diez, C.D. Barr and M. Çetinkaya-Rundel, OpenIntro LaTeX, code, and PDFs are released under a Creative Commons BY-SA 3.0 license.

Question 8.⁵ The expression used in evaluating the *R score* is:

$$R \text{ score} = (Z + ISG + 5) \times 5$$


where *Z* and *ISG* are the numerical expressions for the *Z* score and the indicator of the strength of the group, respectively. The *ISG* is defined as

$$ISG = \frac{\text{average grade of the group at the secondary level} - 75\%}{14}$$

a. (2 marks) Suppose that the average grade of the group at the secondary now taking Statistics and Probability at CEGEP is 86%. The Statistics and Probability class average is 75% and standard deviation is 12%. Compute the R score of a student who has an average of 80%.

b. (2 marks) For a different course, a student earns an R score of 34 what is the student's average in the course if the students taking that course are the same as in part a. and the grades are normally distributed.

c. (2 marks) The following graph is presented on the Marionapolis College website.⁶



Assuming that *Z*-score follow a standard normal distribution and the above claim is true, give a possible explanation of the above results.

⁵CREPUQ. The R score : what it is, and what it does. September 3, 2004

⁶<http://bemarianapolis.ca/choose-us/r-score/>

Bonus Question.⁷ (3 marks)

Bertrand's box paradox is a paradox of elementary probability theory, first posed by Joseph Bertrand in his 1889 work *Calcul des probabilités*.

There are three boxes: a box containing two gold coins, a box containing two silver coins, a box containing one gold coin and a silver coin.

After choosing a box at random and withdrawing one coin at random, if that happens to be a gold coin, what is probability of the next coin also being a gold coin?

⁷Wikipedia contributors. "Bertrand's box paradox." Wikipedia, The Free Encyclopedia. Wikipedia, The Free Encyclopedia, 9 Aug. 2016. Web. 9 Aug. 2016.