

Books, watches, notes or cell phones are not allowed. The only calculators allowed are the Sharp EL-531**. You must show all your work, the correct answer is worth 1 mark the remaining marks are given for the work.

Question 1. (5 marks) Let \mathbf{u} and \mathbf{v} be unit vectors, such that the angle between them is $\frac{2\pi}{3}$. Find $\|5\mathbf{u} - 2\mathbf{v}\|$.

$$\|\mathbf{u}\| = 1$$

$$\|\mathbf{v}\| = 1$$

$$\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \frac{2\pi}{3}$$

$$= 1 \cdot 1 \cdot \left(-\frac{1}{2}\right)$$

$$= -\frac{1}{2}$$

$$\|5\mathbf{u} - 2\mathbf{v}\|^2 = (5\mathbf{u} - 2\mathbf{v}) \cdot (5\mathbf{u} - 2\mathbf{v})$$

$$= (5\mathbf{u}) \cdot (5\mathbf{u}) + (5\mathbf{u}) \cdot (-2\mathbf{v}) + (-2\mathbf{v}) \cdot (5\mathbf{u}) + (-2\mathbf{v}) \cdot (-2\mathbf{v})$$

$$= 25\mathbf{u} \cdot \mathbf{u} - 10\mathbf{u} \cdot \mathbf{v} - 10\mathbf{v} \cdot \mathbf{u} + 4\mathbf{v} \cdot \mathbf{v}$$

$$= 25\|\mathbf{u}\|^2 - 10\mathbf{u} \cdot \mathbf{v} - 10\mathbf{u} \cdot \mathbf{v} + 4\|\mathbf{v}\|^2$$


$$= 25(1)^2 - 20\mathbf{u} \cdot \mathbf{v} + 4(1)^2$$

$$= 29 - 20\left(-\frac{1}{2}\right)$$

$$= 39$$

$$\therefore \|5\mathbf{u} - 2\mathbf{v}\| = \sqrt{39}$$

Question 2. (5 marks) Given a parallelogram with vertices A, B, C and D where $A(2, -2, 4)$ and where the sides of the parallelogram are parallel to $\vec{u} = (1, 2, 3)$ and $\vec{v} = (-3, 2, 5)$. Find the vertices B, C and D .



$$\vec{OB} = \vec{OA} + \vec{u} = (2, -2, 4) + (1, 2, 3) = (3, 0, 7)$$

$$\therefore B(3, 0, 7)$$

$$\vec{OC} = \vec{OA} + \vec{u} + \vec{v} = (2, -2, 4) + (1, 2, 3) + (-3, 2, 5) = (0, 2, 12)$$

$$\therefore C(0, 2, 12)$$

$$\vec{OD} = \vec{OA} + \vec{v} = (2, -2, 4) + (-3, 2, 5) = (-1, 0, 9)$$

$$\therefore D(-1, 0, 9)$$

Question 3. (5 marks) Consider two 4×4 matrices A and B , with $\det(\text{adj}(AB)) = -8$ and $\det(B) = 3$. Find the determinant of M , given

$$\det(\det(B)B^TMA^{-1}) = \det(5\text{adj}(B)A^2).$$

$$(\det(AB))^{4-1} = -8$$

$$(\det B)^4 \det(B^TMA^{-1}) = 5^4 \det(\text{adj}(B)A^2) \quad \det(AB) = -2$$

$$3^4 \det(B^T) \det(M) \det A^{-1} = 5^4 \det(\text{adj}(B)) \det(A^2) \quad \det A \det B = -2$$

$$3^4 \det(B) \det(M) \frac{1}{\det A} = 5^4 (\det B)^{4-1} (\det A)^2 \quad \det A = \frac{-2}{3}$$

$$\det M = \frac{5^4}{3^4} \frac{\det B^3}{\det B} (\det A)^2 \det A$$

$$\det B = \frac{5^4}{3^4} (\det B)^2 (\det A)^3$$

$$\det B = \frac{5^4}{3^{4-2}} \left(\frac{-2}{3}\right)^3$$

$$\det B = -\frac{2^3 5^4}{3^5}$$

Question 4. (3 marks) Determine whether the following statement is true or false. If the statement is false provide a counterexample. If the statement is true provide a proof of the statement.

If A , B and C are $n \times n$ matrices such that $Ax = \mathbf{0}$ has infinitely many solutions then $(A^T C^{-1} + (BA)^T)x = \mathbf{0}$ has infinitely many solutions.

True

By the equivalence thm since $Ax = \mathbf{0}$ has ∞ -many solutions $\Rightarrow \det A = 0$

$$\begin{aligned} \det(A^T C^{-1} + (BA)^T) &= \det(A^T C^{-1} + A^T B) \\ &= \det(A^T (C^{-1} + B)) \\ &= \det A^T \det(C^{-1} + B) \\ &= \det A \det(C^{-1} + B) \\ &= 0 \cdot \det(C^{-1} + B) = 0 \end{aligned}$$

∴ by the equivalence thm. $(A^T C^{-1} + (BA)^T)x = \mathbf{0}$ has infinitely many solutions.