

Integration of Rational Expressions by Partial Fractions

INTRODUCTION:

We start with a few definitions. A **rational expression** is formed when a polynomial is divided by another polynomial. In a **proper rational expression** the degree of the numerator is less than the degree of the denominator. In an **improper rational expression** the degree of the numerator is greater than or equal to the degree of the denominator.

This set of notes is given in three parts. Part A is an explanation of how to decompose a proper rational expression into a sum of simpler fractions. Part B explains Integration by Partial Fractions of proper rational expressions. Part C explains Integration by Partial Fractions of improper rational expressions. Each part includes detailed examples and a set of exercises.

PART A: Partial Fraction Decomposition

In mathematics we often combine two or more rational expressions into one.

$$\text{E.g. } \frac{4}{x+1} + \frac{3}{x-2} = \frac{4(x-2)}{(x+1)(x-2)} + \frac{3(x+1)}{(x-2)(x+1)} = \frac{7x-5}{(x-2)(x+1)}$$

Occasionally, however, the reverse procedure is necessary. The problem is to take a fraction whose denominator is a product of factors, and split it into a sum of simpler fractions. There is more than one way to do this. For the types of expressions we are dealing with the method illustrated here is probably the easiest to apply and understand (no system of equations to solve).

CASE 1 The denominator is a product of distinct linear factors.

For each distinct factor $ax+b$ the sum of partial fractions includes a term of the form $\frac{A}{ax+b}$.

Example 1 Rewrite $\frac{x+5}{(x-4)(x-1)}$ as a sum of simpler fractions.

First write the fraction as $\frac{x+5}{(x-4)(x-1)} = \frac{A}{x-4} + \frac{B}{x-1}$.

Multiply both sides by the common denominator to get $x+5 = A(x-1) + B(x-4)$. In this linear equation we can substitute any x -values to solve for A and B . However, the process is simpler if we choose those values which make a factor zero (i.e. $x-1=0$ or $x-4=0$).

Substitute $x=1$ in the linear equation. $1+5 = A(1-1) + B(1-4) \rightarrow 6 = -3B \rightarrow B = -2$

Substitute $x=4$ in the linear equation. $4+5 = A(4-1) + B(4-4) \rightarrow 9 = 3A \rightarrow A = 3$

Therefore, the partial fraction decomposition is $\frac{x+5}{(x-4)(x-1)} = \frac{3}{x-4} - \frac{2}{x-1}$.

Example 2 Rewrite $\frac{2x-2}{(x+5)(x+2)(x-3)}$ as a sum of simpler fractions.

First write the fraction as $\frac{2x-2}{(x+5)(x+2)(x-3)} = \frac{A}{x+5} + \frac{B}{x+2} + \frac{C}{x-3}$.

Multiply both sides by the common denominator to get the linear equation

$$2x-2 = A(x+2)(x-3) + B(x+5)(x-3) + C(x+5)(x+2)$$

$$\text{Substitute } x=-2. \quad -4-2 = 0+B(-3)(-5)+0 \rightarrow -6 = -15B \rightarrow B = \frac{2}{5}$$

$$\text{Substitute } x=3. \quad 6-2 = 0+0+C(8)(5) \rightarrow 4 = 40C \rightarrow C = \frac{1}{10}$$

$$\text{Substitute } x=-5. \quad -12 = A(-3)(-8)+0+0 \rightarrow -12 = 24A \rightarrow A = -\frac{1}{2}$$

$$\text{The partial fraction decomposition is } \frac{2x-2}{(x+5)(x+2)(x-3)} = -\frac{1}{2} \cdot \frac{1}{x+5} + \frac{2}{5} \cdot \frac{1}{x+2} + \frac{1}{10} \cdot \frac{1}{x-3}$$

CASE 2 The denominator is a product of linear factors, some of which are repeated

For each repeated linear factor $(ax+b)^n$, the sum of partial fractions includes n terms of the

$$\text{form } \frac{A_1}{ax+b} + \frac{A_2}{(ax+b)^2} + \frac{A_3}{(ax+b)^3} + \dots + \frac{A_n}{(ax+b)^n}$$

Example 3 Rewrite $\frac{4x}{(x-2)^2}$ as a sum of simpler fractions.

First write the fraction as $\frac{4x}{(x-2)^2} = \frac{A}{x-2} + \frac{B}{(x-2)^2}$.

Multiply both sides by $(x-2)^2$ to get $4x = A(x-2) + B$.

$$\text{Substitute } x=2. \quad 8 = 0+B \rightarrow B=8$$

There is no other factor to make zero, so we choose an easy x -value to work with.

$$\text{Substitute } x=0 \text{ (and } B=8\text{).} \quad 0 = A(-2)+8 \rightarrow A=4$$

$$\text{The partial fraction decomposition is } \frac{4x}{(x-2)^2} = \frac{4}{x-2} + \frac{8}{(x-2)^2}$$

Example 4 Rewrite $\frac{x^2-2x-5}{x^3-5x^2}$ as a sum of simpler fractions.

(continued next page)

First write the fraction as $\frac{x^2 - 2x - 5}{x^3 - 5x^2} = \frac{x^2 - 2x - 5}{x^2(x-5)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x-5}$.

Multiply by $x^2(x-5)$ to get $x^2 - 2x - 5 = Ax(x-5) + B(x-5) + Cx^2$.

Substitute $x=0$. $-5 = 0 + B(-5) + 0 \rightarrow B = 1$

Substitute $x=5$. $10 = 0 + 0 + C(25) \rightarrow C = \frac{2}{5}$

Substitute $x=1$. $-6 = A(-4) + (-4) + \frac{2}{5}(1) \rightarrow A = \frac{3}{5}$

The partial fraction decomposition is $\frac{x^2 - 2x - 5}{x^3 - 5x^2} = \frac{\frac{3}{5}}{x} + \frac{1}{x^2} + \frac{\frac{2}{5}}{x-5}$.

CASE 3 The denominator has one or more distinct, irreducible quadratic factors.

For each distinct factor $ax^2 + bx + c$ the sum of partial fractions includes a term $\frac{Ax + B}{ax^2 + bx + c}$.

Example 5 Rewrite $\frac{x^2 + 4x + 12}{(x-2)(x^2 + 4)}$ as a sum of simpler fractions.

First write the fraction as $\frac{x^2 + 4x + 12}{(x-2)(x^2 + 4)} = \frac{A}{x-2} + \frac{Bx + C}{x^2 + 4}$.

Multiply by $(x-2)(x^2 + 4)$ to get $x^2 + 4x + 12 = A(x^2 + 4) + (Bx + C)(x-2)$.

Substitute $x=2$. $24 = A(8) + 0 \rightarrow A = 3$

Substitute $x=0$. $12 = 3(4) + (C)(-2) \rightarrow C = 0$

Substitute $x=1$. $17 = 3(5) + (B)(-1) \rightarrow B = -2$

The partial fraction decomposition is $\frac{x^2 + 4x + 12}{(x-2)(x^2 + 4)} = \frac{3}{x-2} - \frac{2x}{x^2 + 4}$.

Example 6 Rewrite $\frac{x^2 + x - 3}{(x+1)(x^2 - 2x + 3)}$ as a sum of simpler fractions.

First write the fraction as $\frac{x^2 + x - 3}{(x+1)(x^2 - 2x + 3)} = \frac{A}{x+1} + \frac{Bx + C}{x^2 - 2x + 3}$.

Multiply by $(x+1)(x^2 - 2x + 3)$ to get $x^2 + x - 3 = A(x^2 - 2x + 3) + (Bx + C)(x+1)$.

Substitute $x=-1$. $-3 = A(6) + 0 \rightarrow A = -\frac{1}{2}$

(continued next page)

Substitute $x=0$ (and $A = -\frac{1}{2}$). $-3 = -\frac{1}{2}(3) + (C)(1) \rightarrow C = -\frac{3}{2}$

Substitute $x=1$ (and $A = -\frac{1}{2}$ and $C = -\frac{3}{2}$). $-1 = -\frac{1}{2}(2) + \left(B - \frac{3}{2}\right)(2) \rightarrow B = \frac{3}{2}$

The partial fraction decomposition is $\frac{x^2+x-3}{(x+1)(x^2-2x+3)} = -\frac{1}{2} \cdot \frac{1}{x+1} + \frac{3}{2} \cdot \frac{x-1}{x^2-2x+3}$.

EXERCISES

Rewrite each of the following as a sum of simpler fractions.

$$1. \frac{4}{(x+1)(x-5)}$$

$$2. \frac{2x-3}{x^2-5x+6}$$

$$3. \frac{5x}{2x^2+11x+12}$$

$$4. \frac{5x+1}{(x+3)(x+2)(x-4)}$$

$$5. \frac{4x^2-x+3}{(x+5)(x-1)(x-2)}$$

$$6. \frac{2x^2-5}{x^3-2x^2-3x}$$

$$7. \frac{x+2}{x^2+8x+16}$$

$$8. \frac{5-4x}{x^3+10x^2+25x}$$

$$9. \frac{5x^2-9x}{(x-4)(x-1)^2}$$

$$10. \frac{x^2-2}{(x+1)(x^2+3)}$$

$$11. \frac{3x^2+9x-4}{(x-1)(x^2+4x-1)}$$

$$12. \frac{6}{(x-5)(x^2-2x+3)}$$

SOLUTIONS

$$1. \frac{4}{(x+1)(x-5)} = \frac{A}{x+1} + \frac{B}{x-5} \rightarrow 4 = A(x-5) + B(x+1) \text{ for all } x$$

$$x=5 \rightarrow B = \frac{2}{3} \quad \text{and} \quad x=-1 \rightarrow A = -\frac{2}{3}$$

$$\frac{4}{(x+1)(x-5)} = \frac{2}{3} \cdot \frac{1}{x-5} - \frac{2}{3} \cdot \frac{1}{x+1}$$

$$2. \frac{2x-3}{x^2-5x+6} = \frac{2x-3}{(x-2)(x-3)} = \frac{A}{x-2} + \frac{B}{x-3} \rightarrow 2x-3 = A(x-3) + B(x-2) \text{ for all } x$$

$$x=3 \rightarrow B = 3 \quad \text{and} \quad x=2 \rightarrow A = -1$$

$$\frac{2x-3}{x^2-5x+6} = \frac{3}{x-3} - \frac{1}{x-2}$$

$$3. \frac{5x}{2x^2+11x+12} = \frac{5x}{(2x+3)(x+4)} = \frac{A}{2x+3} + \frac{B}{x+4} \rightarrow 5x = A(x+4) + B(2x+3)$$

$$x=-4 \rightarrow B=4 \quad \text{and} \quad x=-\frac{3}{2} \rightarrow A=-3$$

$$\frac{5x}{2x^2+11x+12} = \frac{4}{x+4} - \frac{3}{2x+3}$$

$$4. \frac{5x+1}{(x+3)(x+2)(x-4)} = \frac{A}{x+3} + \frac{B}{x+2} + \frac{C}{x-4}$$

$$\rightarrow 5x+1 = A(x+2)(x-4) + B(x+3)(x-4) + C(x+3)(x+2)$$

$$x=-2 \rightarrow B=\frac{3}{2} \quad \text{and} \quad x=4 \rightarrow C=\frac{1}{2} \quad \text{and} \quad x=-3 \rightarrow A=-2$$

$$\frac{5x+1}{(x+3)(x+2)(x-4)} = -\frac{2}{x+3} + \frac{3}{2} \cdot \frac{1}{x+2} + \frac{1}{2} \cdot \frac{1}{x-4}$$

$$5. \frac{4x^2-x+3}{(x+5)(x-1)(x-2)} = \frac{A}{x+5} + \frac{B}{x-1} + \frac{C}{x-2}$$

$$\rightarrow 4x^2-x+3 = A(x-1)(x-2) + B(x+5)(x-2) + C(x+5)(x-1)$$

$$x=1 \rightarrow B=-1 \quad \text{and} \quad x=2 \rightarrow C=\frac{17}{7} \quad \text{and} \quad x=-5 \rightarrow A=\frac{18}{7}$$

$$\frac{4x^2-x+3}{(x+5)(x-1)(x-2)} = \frac{18}{7} \cdot \frac{1}{x+5} - \frac{1}{x-1} + \frac{17}{7} \cdot \frac{1}{x-2}$$

$$6. \frac{2x^2-5}{x^3-2x^2-3x} = \frac{2x^2-5}{x(x+1)(x-3)} = \frac{A}{x} + \frac{B}{x+1} + \frac{C}{x-3}$$

$$\rightarrow 2x^2-5 = A(x+1)(x-3) + Bx(x-3) + Cx(x+1)$$

$$x=-1 \rightarrow B=-\frac{3}{4} \quad \text{and} \quad x=3 \rightarrow C=\frac{13}{12} \quad \text{and} \quad x=0 \rightarrow A=\frac{5}{3}$$

$$\frac{2x^2-5}{x^3-2x^2-3x} = \frac{5}{3} \cdot \frac{1}{x} - \frac{3}{4} \cdot \frac{1}{x+1} + \frac{13}{12} \cdot \frac{1}{x-3}$$

$$7. \frac{x+2}{x^2+8x+16} = \frac{x+2}{(x+4)^2} = \frac{A}{x+4} + \frac{B}{(x+4)^2} \rightarrow x+2 = A(x+4) + B$$

$$x=-4 \rightarrow B=-2 \quad \text{and} \quad x=0, B=-2 \rightarrow A=1$$

$$\frac{x+2}{x^2+8x+16} = \frac{1}{x+4} - \frac{2}{(x+4)^2}$$

$$8. \frac{5-4x}{x^3+10x^2+25x} = \frac{5-4x}{x(x+5)^2} = \frac{A}{x} + \frac{B}{x+5} + \frac{C}{(x+5)^2}$$

$$\rightarrow 5-4x = A(x+5)^2 + Bx(x+5) + Cx$$

$$x=-5 \rightarrow C=-5 \quad \text{and} \quad x=0 \rightarrow A=\frac{1}{5} \quad \text{and} \quad x=1, A=\frac{1}{5}, C=-5 \rightarrow B=-\frac{1}{5}$$

$$\frac{5-4x}{x^3+10x^2+25x} = \frac{1}{5} \cdot \frac{1}{x-3} - \frac{1}{5} \cdot \frac{1}{x+5} - \frac{5}{(x+5)^2}$$

$$9. \frac{5x^2-9x}{(x-4)(x-1)^2} = \frac{A}{x-4} + \frac{B}{x-1} + \frac{C}{(x-1)^2}$$

$$\rightarrow 5x^2-9x = A(x-1)^2 + B(x-4)(x-1) + C(x-4)$$

$$x=1 \rightarrow C=\frac{4}{3} \quad \text{and} \quad x=4 \rightarrow A=\frac{44}{9} \quad \text{and} \quad x=0, A=\frac{44}{9}, C=\frac{4}{3} \rightarrow B=\frac{1}{9}$$

$$\frac{2x^2}{(x-4)(x-1)^2} = \frac{44}{9} \cdot \frac{1}{x-4} + \frac{1}{9} \cdot \frac{1}{x-1} + \frac{4}{3} \cdot \frac{1}{(x-1)^2}$$

$$10. \frac{x^2-2}{(x+1)(x^2+3)} = \frac{A}{x+1} + \frac{Bx+C}{x^2+3} \rightarrow x^2-2 = A(x^2+3) + (Bx+C)(x+1)$$

$$x=-1 \rightarrow A=-\frac{1}{4} \quad \text{and} \quad x=0 \rightarrow C=-\frac{5}{4} \quad \text{and} \quad x=1 \rightarrow B=\frac{5}{4}$$

$$\frac{x^2-2}{(x+1)(x^2+3)} = -\frac{1}{4} \cdot \frac{1}{x+1} + \frac{5}{4} \cdot \frac{x-1}{x^2+3}$$

$$11. \frac{3x^2+9x-4}{(x-1)(x^2+4x-1)} = \frac{A}{x} + \frac{Bx+C}{x^2+4x-1} \rightarrow 3x^2+9x-4 = A(x^2+4x-1) + (Bx+C)(x-1)$$

$$x=1 \rightarrow A=2 \quad \text{and} \quad x=0 \rightarrow C=2 \quad \text{and} \quad x=-1 \rightarrow B=1$$

$$\frac{3x^2+9x-4}{(x-1)(x^2+4x-1)} = \frac{2}{x-1} + \frac{x+2}{x^2+4x-1}$$

$$12. \frac{6}{(x-5)(x^2-2x+3)} = \frac{A}{x-5} + \frac{Bx+C}{x^2-2x+3} \rightarrow 6 = A(x^2-2x+3) + (Bx+C)(x-5)$$

$$x=5 \rightarrow A=\frac{1}{3} \quad \text{and} \quad x=0 \rightarrow C=-1 \quad \text{and} \quad x=1 \rightarrow B=-\frac{1}{3}$$

$$\frac{6}{(x-5)(x^2-2x+3)} = \frac{1}{3} \cdot \frac{1}{x-5} - \frac{1}{3} \cdot \frac{x+3}{x^2-2x+3}$$

PART B:

Integration of proper Rational Expressions by Partial Fractions

In this part the student is expected to understand partial fraction decomposition as explained in Part A. The student is also expected to be able to perform elementary integrations (by substitution or by inspection) of the following types.

$$\int \frac{1}{ax+b} dx = \frac{1}{a} \cdot \ln|ax+b| + C$$

$$\int \frac{1}{(ax+b)^2} dx = -\frac{1}{a} \cdot \frac{1}{ax+b} + C$$

$$\int \frac{2ax+b}{ax^2+bx+c} dx = \ln|ax^2+bx+c| + C$$

In Part B each indefinite integral (antiderivative) must be simplified by decomposing the proper rational expression into a sum of partial fractions. The details of the partial fraction decomposition are left to the student. If necessary go back and review Part A.

Example 1 Find the indefinite integral. $\int \frac{7x+1}{(x+3)(x-1)} dx$

$$\int \frac{7x+1}{(x+3)(x-1)} dx = \int \left[\frac{5}{x+3} + \frac{2}{x-1} \right] dx = 5 \ln|x+3| + 2 \ln|x-1| + C$$

Example 2 Find the indefinite integral. $\int \frac{6x^2+2x}{(x+3)(x-1)(x-5)} dx$

$$\int \frac{6x^2+2x}{(x+3)(x-1)(x-5)} dx = \int \left[\frac{3}{2} \cdot \frac{1}{x+3} - \frac{1}{2} \cdot \frac{1}{x-1} + \frac{5}{x-5} \right] dx = \frac{3}{2} \ln|x-3| - \frac{1}{2} \ln|x-1| + 5 \ln|x-5| + C$$

Example 3 Find the indefinite integral. $\int \frac{1-3x}{x^3+4x^2+4x} dx$

$$\int \frac{1-3x}{x(x+2)^2} dx = \int \left[\frac{1}{4} \cdot \frac{1}{x} - \frac{1}{4} \cdot \frac{1}{x+2} - \frac{7}{2} \cdot \frac{1}{(x+2)^2} \right] dx = \frac{1}{4} \ln|x| - \frac{1}{4} \ln|x+2| + \frac{7}{2} \cdot \frac{1}{x+2} + C$$

Example 4 Find the indefinite integral. $\int \frac{x-3}{(x+2)(x^2+6)} dx$

(continued next page)

$$\int \frac{x-3}{(x+2)(x^2+6)} dx = \int \left[-\frac{1}{2} \cdot \frac{1}{x+2} + \frac{1}{2} \cdot \frac{x}{x^2+6} \right] dx = -\frac{1}{2} \ln|x+2| + \frac{1}{4} \ln|x^2+6| + C$$

Example 5 Find the indefinite integral.

$$\int \frac{x^2+4x-7}{(x-3)(x^2+6x-6)} dx = \int \left[\frac{2}{3} \cdot \frac{1}{x-3} + \frac{1}{3} \cdot \frac{x+3}{x^2+6x-6} \right] dx = \frac{2}{3} \ln|x-3| + \frac{1}{6} \ln|x^2+6x-6| + C$$

EXERCISES Find the indefinite integral.

1. $\int \frac{3-4x}{x^2+x} dx$

2. $\int \frac{x}{x^2+7+10} dx$

3. $\int \frac{6}{3x^2-14x+8} dx$

4. $\int \frac{3x^2+8x-7}{(x+4)(x+3)(x+1)} dx$

5. $\int \frac{2-4x^2}{(x+2)(x-2)(x-5)} dx$

6. $\int \frac{3x}{(x+4)(x-1)(x-3)} dx$

7. $\int \frac{3-2x}{x^2+6x+9} dx$

8. $\int \frac{3x-1}{x^3-2x^2} dx$

9. $\int \frac{2x^2+x+4}{(x+1)(x-4)^2} dx$

10. $\int \frac{5x^2+8x+6}{(x+4)(x^2+2)} dx$

11. $\int \frac{12x+18}{(x+3)(2x^2+8x+9)} dx$

12. $\int \frac{15-25x}{(x-4)(2x^2-6x+9)} dx$

ANSWERS

1. $\int \frac{3-4x}{x(x+1)} dx = \int \left[\frac{3}{x} - \frac{7}{x+1} \right] dx = 3 \ln|x| - 7 \ln|x+1| + C$

2. $\int \frac{x}{(x+5)(x+2)} dx = \int \left[\frac{5}{3} \cdot \frac{1}{x+5} - \frac{2}{3} \cdot \frac{1}{x+2} \right] dx = \frac{5}{3} \cdot \ln|x+5| - \frac{2}{3} \cdot \ln|x+2| + C$

3. $\int \frac{6}{(3x-2)(x-4)} dx = \int \left[-\frac{9}{5} \cdot \frac{1}{3x-2} + \frac{3}{5} \cdot \frac{1}{x-4} \right] dx = -\frac{3}{5} \cdot \ln|3x-2| + \frac{3}{5} \cdot \ln|x-4| + C$

4. $\int \left[\frac{3}{x+4} + \frac{2}{x+3} - \frac{2}{x+1} \right] dx = 3 \ln|x+4| + 2 \ln|x+3| - 2 \ln|x+1| + C$

$$5. \int \left[-\frac{1}{2} \cdot \frac{1}{x+2} + \frac{7}{6} \cdot \frac{1}{x-2} - \frac{14}{3} \cdot \frac{1}{x-5} \right] dx = -\frac{1}{2} \cdot \ln|x+2| + \frac{7}{6} \cdot \ln|x-2| - \frac{14}{3} \cdot \ln|x-5| + C$$

$$6. \int \left[-\frac{12}{35} \cdot \frac{1}{x+4} - \frac{3}{10} \cdot \frac{1}{x-1} + \frac{9}{14} \cdot \frac{1}{x-3} \right] dx = -\frac{12}{35} \cdot \ln|x+4| - \frac{3}{10} \cdot \ln|x-1| + \frac{9}{14} \cdot \ln|x-3| + C$$

$$7. \int \frac{3-2x}{(x+3)^2} dx = \int \left[\frac{-2}{x+3} + \frac{9}{(x+3)^2} \right] dx = -2 \ln|3x+2| - \frac{9}{x+3} + C$$

$$8. \int \frac{3x-1}{x^2(x-2)} dx = \int \left[-\frac{5}{4} \cdot \frac{1}{x} + \frac{1}{2} \cdot \frac{1}{x^2} + \frac{5}{4} \cdot \frac{1}{x-2} \right] dx = -\frac{5}{4} \cdot \ln|x| - \frac{1}{2} \cdot \frac{1}{x} + \frac{5}{4} \cdot \ln|x-2| + C$$

$$9. \int \left[\frac{1}{5} \cdot \frac{1}{x+1} + \frac{9}{5} \cdot \frac{1}{x-4} + \frac{8}{(x-4)^2} \right] dx = \frac{1}{5} \cdot \ln|x+1| + \frac{9}{5} \cdot \ln|x-4| - \frac{8}{x-4} + C$$

$$10. \int \left[\frac{3}{x+4} + \frac{2x}{x^2+2} \right] dx = 3 \ln|x+4| + \ln|x^2+2| + C$$

$$11. \int \left[\frac{-6}{x+3} + \frac{12x+24}{2x^2+8x+9} \right] dx = -6 \ln|x+3| + 3 \ln|2x^2+8x+9| + C$$

$$12. \int \left[\frac{-5}{x-4} + \frac{10x-15}{2x^2-6x+9} \right] dx = -5 \ln|x-4| + \frac{5}{2} \ln|2x^2-6x+9| + C$$

PART C:

Integration of improper Rational Expressions by Partial Fractions

The prerequisite skills of Part B are also required in Part C. Furthermore, the student is expected to be able to use long division to decompose an improper rational expression.

In each indefinite integral of Part C the improper rational expression must be rewritten as a polynomial plus proper rational. In most cases the resulting proper rational can then be further simplified by decomposition into a sum of partial fractions. The details of the long division and partial fraction decomposition are left to the student.

Example 1 Find the indefinite integral. $\int \frac{3x^3+8x^2-10x+15}{x+4} dx$

$$\int \frac{3x^3+8x^2-10x+15}{x+4} dx = \int \left[3x^2 - 4x + 6 - \frac{9}{x+4} \right] dx = x^3 - 2x^2 + 6x - 9 \ln|x+4| + C$$

Example 2 Find the indefinite integral.

$$\int \frac{2x^2+4x-7}{x^2+x-6} dx = \int \left[2 + \frac{2x+5}{(x+3)(x-2)} \right] dx = \int \left[2 + \frac{1}{5} \cdot \frac{1}{x+3} + \frac{9}{5} \cdot \frac{1}{x-2} \right] dx$$

$$\rightarrow \int \frac{2x^2+4x-7}{x^2+x-6} dx = 2x + \frac{1}{5} \cdot \ln|x+3| + \frac{9}{5} \cdot \ln|x-2| + C$$

Example 3 Find the indefinite integral.

$$\int \frac{2x^3+2x^2-95x+40}{x^3+x^2-20x} dx = \int \left[2 + \frac{40-11x}{x(x+5)(x-4)} \right] dx = \int \left[2 - \frac{2}{x} + \frac{7}{x+5} - \frac{5}{x-4} \right] dx$$

$$\rightarrow \int \frac{2x^3+2x^2-95x+40}{x^3+x^2-20x} dx = 2x - 2 \ln|x| + 7 \ln|x+5| - 5 \ln|x-4| + C$$

Example 4 Find the indefinite integral.

$$\int \frac{2x^4-9x^3+9x^2-8x-6}{x^3-6x^2+9x} dx = \int \left[2x + 3 + \frac{9x^2-35x-6}{x(x-3)^2} \right] dx$$

$$= \int \left[2x + 3 - \frac{2}{3} \cdot \frac{1}{x} + \frac{29}{3} \cdot \frac{1}{x-3} - \frac{10}{(x-3)^2} \right] dx$$

$$= x^2 + 3x - \frac{2}{3} \cdot \ln|x| + \frac{29}{3} \cdot \ln|x-3| + \frac{10}{x-3} + C$$

EXERCISES Find the indefinite integral.

1. $\int \frac{6x+5}{x+2} dx$

2. $\int \frac{4x^2-12x-25}{x-5} dx$

3. $\int \frac{5x^3+3x-2}{x-1} dx$

4. $\int \frac{x^3+3x^2-4x-6}{x^2+2x-15} dx$

5. $\int \frac{4x^2-8x+3}{x^2-3x-4} dx$

6. $\int \frac{x^3-3x^2}{x^2-3x-10} dx$

7. $\int \frac{4x^3+20x^2+15x+8}{x^3+5x^2+4x} dx$

8. $\int \frac{x^4-10x^3+28x^2-15x-15}{x^3-7x^2+10x} dx$

$$9. \int \frac{x^5 - 2x^4 - 7x^3 + 20x^2 - 12x + 4}{x^3 + x^2 - 6x} dx$$

$$10. \int \frac{4x^5 + 6x^4 + 2x^3 + 3x^2 - 5x - 7}{x^3 + 2x^2 + x} dx$$

$$11. \int \frac{2x^3 - 15x^2 + 17x + 25}{x^3 - 10x^2 + 25x} dx$$

$$12. \int \frac{x^4 + 2x^3 - 6x^2 - 6x + 3}{x^3 + 4x^2 + 4x} dx$$

ANSWERS

$$1. \int \frac{6x+5}{x+2} dx = \int \left[6 - \frac{7}{x+2} \right] dx = 6x - 7 \ln|x+2| + C$$

$$2. \int \frac{4x^2 - 12x - 25}{x-5} dx = \int \left[4x + 8 + \frac{15}{x-5} \right] dx = 2x^2 + 8x + 15 \ln|x-5| + C$$

$$3. \int \frac{5x^3 + 3x - 2}{x-1} dx = \int \left[5x^2 + 5x + 8 + \frac{6}{x-1} \right] dx = \frac{5}{3}x^3 + \frac{5}{2}x^2 + 8x + 6 \ln|x-1| + C$$

$$4. \int \frac{x^3 + 3x^2 - 4x - 6}{x^2 + 2x - 15} dx = \int \left[x + 1 + \frac{9x+9}{(x+5)(x-3)} \right] dx \\ = \int \left[x + 1 + \frac{9}{2} \cdot \frac{1}{x+5} + \frac{9}{2} \cdot \frac{1}{x-3} \right] dx = \frac{1}{2}x^2 + x + \frac{9}{2} \cdot \ln|x+5| + \frac{9}{2} \cdot \ln|x-3| + C$$

$$5. \int \frac{4x^2 - 8x + 3}{x^2 - 3x - 4} dx = \int \left[4 + \frac{4x+19}{(x+1)(x-4)} \right] dx = \int \left[4 - \frac{3}{x+1} + \frac{7}{x-4} \right] dx \\ = 4x - 3 \ln|x+1| + 7 \ln|x-4| + C$$

$$6. \int \frac{x^3 - 3x^2}{x^2 - 3x - 10} dx = \int \left[x + \frac{10x}{(x+2)(x-5)} \right] dx = \int \left[x + \frac{20}{7} \cdot \frac{1}{x+2} + \frac{50}{7} \cdot \frac{1}{x-5} \right] dx \\ = \frac{1}{2}x^2 + \frac{20}{7} \cdot \ln|x+2| + \frac{50}{7} \cdot \ln|x-5| + C$$

$$7. \int \frac{4x^3 + 20x^2 + 15x + 8}{x^3 + 5x^2 + 4x} dx = \int \left[4 + \frac{8-x}{x(x+4)(x+1)} \right] dx = \int \left[4 + \frac{2}{x} + \frac{1}{x+4} - \frac{3}{x+1} \right] dx \\ = 4x + 2 \ln|x| + \ln|x+4| - 3 \ln|x+1| + C$$

$$8. \int \frac{x^4 - 10x^3 + 28x^2 - 15x - 15}{x^3 - 7x^2 + 10x} dx = \int \left[x - 3 + \frac{-3x^2 + 15x - 15}{x(x-2)(x-5)} \right] dx \\ = \int \left[x - 3 - \frac{3}{2} \cdot \frac{1}{x} - \frac{1}{2} \cdot \frac{1}{x-2} - \frac{1}{x-5} \right] dx = \frac{1}{2}x^2 - 3x - \frac{3}{2}\ln|x| - \frac{1}{2}\ln|x-2| - \ln|x-5| + C$$

$$9. \int \frac{x^5 - 2x^4 - 7x^3 + 20x^2 - 12x + 4}{x^3 + x^2 - 6x} dx = \int \left[x^2 - 3x + 2 + \frac{4}{x(x+3)(x-2)} \right] dx \\ = \int \left[x^2 - 3x + 2 - \frac{2}{3} \cdot \frac{1}{x} + \frac{4}{15} \cdot \frac{1}{x+3} + \frac{2}{5} \cdot \frac{1}{x-2} \right] dx \\ = \frac{1}{3}x^3 - \frac{3}{2}x^2 + 2x - \frac{2}{3}\ln|x| + \frac{4}{15}\ln|x+3| + \frac{2}{5}\ln|x-2| + C$$

$$10. \int \frac{4x^5 + 6x^4 + 2x^3 + 3x^2 - 5x - 7}{x^3 + 2x^2 + x} dx = \int \left[4x^2 - 2x + 2 + \frac{x^2 - 7x - 7}{x(x+1)^2} \right] dx \\ = \int \left[4x^2 - 2x + 2 - \frac{7}{x} + \frac{8}{x+1} - \frac{1}{(x+1)^2} \right] dx \\ = \frac{4}{3}x^3 - x^2 + 2x - 7\ln|x| + 8\ln|x+1| + \frac{1}{x+1} + C$$

$$11. \int \frac{2x^3 - 15x^2 + 17x + 25}{x^3 - 10x^2 + 25x} dx = \int \left[2 + \frac{5x^2 - 33x + 25}{x(x-5)^2} \right] dx = \int \left[2 + \frac{1}{x} + \frac{4}{x-5} - \frac{3}{(x-5)^2} \right] dx \\ = 2x + \ln|x| + 4\ln|x-5| - \frac{3}{x-5} + C$$

$$12. \int \frac{x^4 + 2x^3 - 6x^2 - 6x + 3}{x^3 + 4x^2 + 4x} dx = \int \left[x - 2 + \frac{-2x^2 + 2x + 3}{x(x+2)^2} \right] dx \\ = \int \left[x - 2 + \frac{3}{4} \cdot \frac{1}{x} - \frac{11}{4} \cdot \frac{1}{x+2} + \frac{9}{2} \cdot \frac{1}{(x+2)^2} \right] dx \\ = \frac{1}{2}x^2 - 2x + \frac{3}{4}\ln|x| - \frac{11}{4}\ln|x+1| - \frac{9}{2} \cdot \frac{1}{x+2} + C$$