Name:
 \_\_\_\_\_\_

 Student ID:
 \_\_\_\_\_\_

## Test 2

This test is graded out of 50 marks. No books, notes, graphing calculators or cell phones are allowed. You must show all your work, the correct answer is worth 1 mark the remaining marks are given for the work. If you need more space for your answer use the back of the page.

**Question 1.** (5 marks) Use Cramer's rule to solve for  $x_2$  without solving for  $x_1$ ,  $x_3$ .

$3x_1$	+	$2x_2$	+	$2x_3$	=	1
$4x_1$	+	$x_2$			=	0
$7x_1$	+	$3x_2$	_	$x_3$	=	3

(Use cofactor expansions to find the determinants)

**Question 2.** (4 marks) Show that if A, B are invertible then  $det(A^{-1}BCAB^{-1}) = det(C)$ 

## Question 3. (5 marks)

a. (5 marks) Use the combinatorial definition of the determinant to compute:

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

b. (2 marks) Justify the "visual way" of computing the determinant using part a.

## **Question 4.**

a. (5 marks) Find the inverse of the following matrix using the adjoint:

$$A = \begin{bmatrix} 1 & -2 \\ 3 & 0 \end{bmatrix}$$

b. (2 marks) Using part a. solve the following equation:

$$Ax = b$$

where

$$b = \begin{bmatrix} 1 \\ 3 \end{bmatrix}.$$

**Question 5.** (3 marks) For which value(s) of  $\alpha$  is the following matrix invertible:

α	-2	2 ]
0	$\alpha - 1$	101
0	0	$\alpha^2 - 1$

Question 6. Let

A =	0 0 0 0 0 0 0 0 0	1 0 0 0 0 0 0 0 0	1 0 0 0 0 0 0 0 0	1 1 0 0 0 0 0 0 0	9 4 1 0 0 0 0 0 0	3 9 4 1 1 0 0 0 0	4 9 4 1 1 0 0 0	1 7 9 4 1 1 0 0	0 7 2 9 4 1 1 0	9 7 7 2 9 4 1 1	$, C = \begin{bmatrix} a \\ c \end{bmatrix}$	$\begin{bmatrix} b \\ d \end{bmatrix}, D = \begin{bmatrix} \end{bmatrix}$	$\begin{bmatrix} b & 3a-2b \\ d & 3c-2d \end{bmatrix}$
	0	0	0	0	0	0	0	0	0	1			
	0	0	0	0	0	0	0	0	0	0			

a. (3 marks) If B is a  $10 \times 10$  matrix show that AB is not invertible.

b. (4 marks) If det(D) = 2 then find det(C).

Question 7. (5 marks) Compute the determinant of the following matrix using elementary operations

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 3 & -2 & 1 \\ 4 & 3 & 1 \end{bmatrix}.$$

**Question 8.** (5 marks) If A, B are  $3 \times 3$  matrices, det(2A) = -8 and det $(B) = \sqrt{2}$  then find

 $\det((3AB)^{-1}(2AB)^t A^3 B^4 B^{-1}).$ 

(show every step)

## Question 9.

- a. (1 mark) Sketch the vector  $\mathbf{v} = (2,3,5)$  on a right-handed coordinate system.
- b. (2 marks) Find a nonzero vector **u** with terminal point Q(1,2,3) which is oppositely directed to **v**.
- c. (2 marks) Let  $\mathbf{a} = (1, 0, -2)$ ,  $\mathbf{b} = (0, 2, 0)$  and  $\mathbf{c} = (0, 1, 1)$ . If  $\mathbf{w} = 2(\mathbf{a} \mathbf{b}) + 3\mathbf{c}$  then find a unit vector which has the same direction as  $\mathbf{w}$

Bonus Question. (3 marks) Consider

$$A = \begin{bmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{bmatrix}$$

and use row reduction to show that det(A) = (b-a)(c-a)(c-b).