

No books, watches, notes or cell phones are allowed. The only calculators allowed are the Sharp EL-531**. You must show all your work, the correct answer is worth 1 mark the remaining marks are given for the work.

Question 1.¹ (1 mark per blank) Complete each of the following sentences with MUST, MIGHT, or CANNOT.

- If A is a product of elementary matrices, then $\det(A)$ might equal one.
- Let A be a 3×3 matrix, and let B be a 4×4 matrix. If the leading ones of the RREF of A is equal to those of the RREF of B , then $\det(A)$ might equal zero and $\det(B)$ must equal zero.
- If \vec{u} and \vec{v} are nonzero vectors and $\text{proj}_{\vec{v}} \vec{u} = \vec{u}$, then \vec{u} must be parallel to \vec{v} .
- Let \vec{w} be orthogonal to both \vec{u} and \vec{v} . Then \vec{w} must be orthogonal to $\vec{u} + \vec{v}$.
- The vector $\vec{u} \times (\vec{v} \times \vec{w})$ must be parallel to the vector \vec{u} .
- The vector $\vec{u} \times (\vec{v} \times \vec{w})$ cannot be orthogonal to the vector $2\vec{v} \times (-4\vec{w})$.

Question 2. (1 mark per blank) Given A an $n \times n$ matrix and k a non-zero scalar.

- If A is the reduced row echelon form of a singular matrix then $\det(A) = \underline{0}$.
- If A is an elementary matrix obtained by adding k times one row to another then $\det(A) = \underline{1}$.
- If A is the identity matrix multiplied by k then $\det(A) = \underline{k^n}$.

Question 3. (5 marks)² Given A , an $n \times n$ matrix such that $\det(A) = 9$ and

$$A^3 A^T = 3A^{-1} \text{adj}(A)$$

find n .

$$\det(A^3 A^T) = \det(3A^{-1} \text{adj}(A))$$

$$\det(A^3) \det(A^T) = 3^n \det(A^{-1}) \det(\text{adj}(A))$$

$$(\det(A))^3 \det A = 3^n \frac{1}{\det(A)} (\det(A))^{n-1}$$

$$(\det(A))^5 = 3^n (9)^{n-1}$$

$$9^5 = 3^n (3^2)^{n-1}$$

$$(3^2)^5 = 3^n 3^{2(n-1)}$$

$$3^{10} = 3^{n+2n-2}$$

$$10 = n+2n-2$$

$$12 = 3n$$

$$4 = n$$

Question 4. (5 marks) Using elementary operations show that

$$-sr \begin{vmatrix} a & b \\ c & d \end{vmatrix} = \begin{vmatrix} sb+2d & rsa+2rc \\ d & rc \end{vmatrix} \xrightarrow{c_1 \leftrightarrow c_2} -sr \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

$$\text{RHS} = \begin{vmatrix} sb+2d & rsa+2rc \\ d & rc \end{vmatrix}$$

$$= -2R_2 + R_1 \rightarrow R_1 \begin{vmatrix} sb & rsa \\ d & rc \end{vmatrix}$$

$$= \frac{1}{s} R_1 \rightarrow R_1 \begin{vmatrix} b & ra \\ d & rc \end{vmatrix}$$

$$= \frac{1}{r} C_2 \rightarrow C_2 \begin{vmatrix} b & a \\ d & c \end{vmatrix}$$

¹ From or modified from a John Abbott final examination

² From a Dawson College final examination

Question 4. Given $A^{-1} = \begin{bmatrix} 1 & 0 & 2 & 3 \\ -1 & 0 & 0 & 1 \\ 4 & -2 & 0 & 0 \\ 0 & 0 & 5 & 1 \end{bmatrix}$.

a. (5 marks) Find $\det(A)$.

b. (2 marks) Find $\text{adj}(A)$.

$$A^{-1} = \frac{1}{\det(A)} \text{adj}(A)$$

$$\text{adj}(A) = \det(A) A^{-1}$$

$$= \frac{1}{-36} \begin{bmatrix} 1 & 0 & 2 & 3 \\ -1 & 0 & 0 & 1 \\ 4 & -2 & 0 & 0 \\ 0 & 0 & 5 & 1 \end{bmatrix}$$

$$\begin{aligned} |A^{-1}| &= \underbrace{a_{12}C_{12}}_0 + \underbrace{a_{22}C_{22}}_0 + a_{32}C_{32} + \underbrace{a_{42}C_{42}}_0 \\ &= a_{32}C_{32} \\ &= -2(-1)^{3+2} \begin{vmatrix} 1 & 2 & 3 \\ -1 & 0 & 1 \\ 0 & 5 & 1 \end{vmatrix} \end{aligned}$$

$$= 2 \begin{vmatrix} 1 & 2 & 3 \\ 0 & 2 & 4 \\ 0 & 5 & 1 \end{vmatrix} \text{ after } R_1 + R_2 \rightarrow R_2$$

$$= 2 [a_{11}C_{11} + a_{21}C_{21} + a_{31}C_{31}]$$

$$= 2(1)(-1)^{1+1} \begin{vmatrix} 2 & 4 \\ 5 & 1 \end{vmatrix} = 2(-16) = -36$$

$$\det(A) = \frac{1}{\det(A^{-1})} = \frac{1}{-36}$$

Question 5.¹ (3 marks) Determine whether the following statement is true or false. If the statement is false provide a counterexample. If the statement is true provide a proof of the statement.

If A is a symmetric $n \times n$ matrix where n is even then $\det(A) = 0$.

False,

$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ is symmetric
since $I_2^T = I_2$, $n=2$ (even)
but $\det(I_2) = 1$.

Question 6. (3 marks) Solve only for x_3 using Cramer's Rule.


$$\begin{cases} x_1 - 2x_2 + 3x_3 = 4 \\ 5x_2 - 6x_3 = 7 \\ 8x_3 = 9 \end{cases} \quad \underbrace{\begin{bmatrix} 1 & -2 & 3 \\ 0 & 5 & -6 \\ 0 & 0 & 8 \end{bmatrix}}_A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \underbrace{\begin{bmatrix} 4 \\ 7 \\ 9 \end{bmatrix}}_b$$

$$\det A = 1 \cdot 5 \cdot 8$$

$$\det A_3 = \det \begin{bmatrix} 1 & -2 & 4 \\ 0 & 5 & 7 \\ 0 & 0 & 9 \end{bmatrix} = 1 \cdot 5 \cdot 9$$

$$x_3 = \frac{\det A_3}{\det A} = \frac{1 \cdot 5 \cdot 9}{1 \cdot 5 \cdot 8} = \frac{9}{8}$$

Question 7. (2 marks) Using sketches illustrate that vector addition is commutative.



Question 8.¹ Let \vec{u} and \vec{v} be vectors in \mathbb{R}^n . Given: $\|\vec{u}\| = 5$, $\|\vec{u} + 2\vec{v}\| = \sqrt{2}$, \vec{v} and $\vec{u} + 3\vec{v}$ are both unit vectors, and the angle between $\vec{u} + 2\vec{v}$ and $\vec{u} + 3\vec{v}$ is $\pi/4$.

- (3 marks) Find $\vec{u} \cdot \vec{v}$.
- (2 marks) Find $\|\vec{u} + \vec{v}\|$.

$$\begin{aligned} a) (\vec{u} + 2\vec{v}) \cdot (\vec{u} + 3\vec{v}) &= \|\vec{u} + 2\vec{v}\| \|\vec{u} + 3\vec{v}\| \cos \frac{\pi}{4} \\ \vec{u} \cdot \vec{u} + \vec{u} \cdot (3\vec{v}) + (2\vec{v}) \cdot \vec{u} + (2\vec{v}) \cdot (3\vec{v}) &= \sqrt{2} \cdot 1 \cdot \frac{1}{\sqrt{2}} \\ \|\vec{u}\|^2 + 3\vec{u} \cdot \vec{v} + 2\vec{v} \cdot \vec{u} + 6\|\vec{v}\|^2 &= 1 \\ 5^2 + 3\vec{u} \cdot \vec{v} + 2\vec{u} \cdot \vec{v} + 6\|\vec{v}\|^2 &= 1 \\ 5\vec{u} \cdot \vec{v} &= 1 - 25 - 6(1^2) \\ \vec{u} \cdot \vec{v} &= -6 \end{aligned}$$

$$\begin{aligned} b) \|\vec{u} + \vec{v}\|^2 &= (\vec{u} + \vec{v}) \cdot (\vec{u} + \vec{v}) \\ &= \vec{u} \cdot \vec{u} + \vec{u} \cdot \vec{v} + \vec{v} \cdot \vec{u} + \vec{v} \cdot \vec{v} \\ &= \|\vec{u}\|^2 + 2\vec{u} \cdot \vec{v} + \|\vec{v}\|^2 \\ &= 5^2 + 2(-6) + 1^2 \\ &= 14 \end{aligned}$$

Question 9. Given two planes: $\mathcal{P}_1 : x + z = 1$ and $\mathcal{P}_2 : y + z = 1$.

- (1 mark) Give a point of intersection of the two planes, by inspection.

$(0, 0, 1)$ satisfies both \mathcal{P}_1 and \mathcal{P}_2 .

- (1 mark) Give a geometrical argument to explain why the intersection of the two planes is a line.

The normals of both planes are not parallel. Hence the inclination of both planes are different. \therefore the intersection of \mathcal{P}_1 and \mathcal{P}_2 is a line.


- (2 marks) Find a direction vector for the line of intersection of the two planes without solving for the solution set.

The intersection is parallel to both planes, therefore orthogonal to both normals. $\vec{d} = \vec{n}_1 \times \vec{n}_2 = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} = (-1, -1, 1)$

- (2 marks) Find the solution set of the system of linear equations determined by \mathcal{P}_1 and \mathcal{P}_2 by only using part a) and part c).

Justify. $\vec{x} =$ particular solution of $A\vec{x} = \vec{b}$ + general solution of $A\vec{x} = \vec{0}$
 $= (0, 0, 1) + t(-1, -1, 1)$ where $t \in \mathbb{R}$ since the general solution of $A\vec{x} = \vec{0}$ is the line which passes through the origin and $\perp \vec{n}_1, \vec{n}_2$

- (1 marks) Sketch a geometrical interpretation to part d).




Question 10. Given the points $A(3, 0, -1)$ and $B(3, 1, 0)$.

a. (1 mark) Find the equation of the line which passes through A and B .

$\mathcal{L}: \underline{x} = \underline{A} + t\underline{AB} \quad t \in \mathbb{R} \quad \underline{AB} = \underline{B} - \underline{A} = (3, 1, 0) - (3, 0, -1) = (0, 1, 1)$

b. (4 marks) Find the points on the line which passes through A and B which are $\sqrt{21}$ units away from the origin.



$\mathcal{L}: \underline{x} = (3, 0, -1) + t(0, 1, 1) = (3, t, -1+t)$

$\|Ox\| = \sqrt{21}$
 $\|x\| = \sqrt{21}$
 $\sqrt{21} = \sqrt{3^2 + t^2 + (-1+t)^2}$
 $21 = 9 + t^2 + 1 - 2t + t^2$
 $0 = 2t^2 - 2t - 11$

\therefore Two points, when $t_1 = \frac{1+\sqrt{23}}{2}$

and $t_2 = \frac{1-\sqrt{23}}{2}$

$P_1 = (3, 0, -1) + t_1(0, 1, 1)$

$P_2 = (3, 0, -1) + t_2(0, 1, 1)$

So $t = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(2)(-11)}}{2(2)} = \frac{2 \pm \sqrt{4+88}}{4} = \frac{2 \pm \sqrt{92}}{4} = \frac{2 \pm 2\sqrt{23}}{4} = \frac{1 \pm \sqrt{23}}{2}$

Question 11.³ (4 marks) Determine whether the two lines $\mathcal{L}_1: \underline{x} = (1 - 2t, 2 - t, 3 + 3t)$ and $\mathcal{L}_2: \underline{x} = (4 + 3t, -1 + t, 2 + t)$ intersect, are parallel or are skew lines.

$\mathcal{L}_1: \underline{x} = (1, 2, 3) + t(-2, -1, 3), t \in \mathbb{R}$
 \underline{d}_1

$\mathcal{L}_2: \underline{x} = (4, -1, 2) + s(3, 1, 1), s \in \mathbb{R}$
 \underline{d}_2

\mathcal{L}_1 and \mathcal{L}_2 are not \parallel since $\underline{d}_1 \nparallel \underline{d}_2$, because they are not multiples of each other.

Do \mathcal{L}_1 and \mathcal{L}_2 intersect?

$\begin{cases} 1-2t = 4+3s \\ 2-t = -1+s \\ 3+3t = 2+s \end{cases} \Rightarrow \begin{cases} -3s-2t = 3 \\ -s-t = -3 \\ -s+3t = -1 \end{cases}$

$\begin{bmatrix} -1 & -1 & -3 \\ -1 & 3 & -1 \\ -3 & -2 & 3 \end{bmatrix}$


$\sim \begin{matrix} -R_1 + R_2 \rightarrow R_2 \\ -3R_1 + R_3 \rightarrow R_3 \end{matrix} \begin{bmatrix} -1 & -1 & -3 \\ 0 & 4 & 2 \\ 0 & 1 & 12 \end{bmatrix}$

$\sim \begin{matrix} -\frac{1}{4}R_2 + R_3 \rightarrow R_3 \end{matrix} \begin{bmatrix} -1 & -1 & -3 \\ 0 & 4 & 2 \\ 0 & 0 & \frac{23}{2} \end{bmatrix}$

\therefore inconsistent
 $\therefore \nexists s, t$, for which the lines have the same point. $\therefore \mathcal{L}_1$ and \mathcal{L}_2 do not intersect. $\therefore \mathcal{L}_1$ and \mathcal{L}_2 are skew lines.

Question 12. (5 marks) A cylinder where the sides are not perpendicular to the base (the circle) is called an *oblique cylinder*. Given the oblique cylinder defined by the given vectors $\vec{u} = (2, 2, 4)$, $\vec{v} = (1, 2, 1)$ and $\vec{w} = (4, 1, 3)$. Find the volume of the oblique cylinder.

Note that from the diagram that \vec{w} is not perpendicular to the base, that \vec{v} is positioned such that its tail is at the center of the circle and its tip lies on the circle, that \vec{u} is positioned such that the vector passes through the center of the circle while its tail and tip lie on the circle. (Hint: the volume of an oblique cylinder is equal to the area of the base times the height.)



Area of base = $\pi r^2 = \pi (\|\vec{v}\|)^2 = \pi (\sqrt{(1)^2 + (2)^2 + (1)^2})^2 = 6\pi$

height = $\|\text{proj}_{\vec{v}} \vec{u}\|$

$\vec{v} \times \vec{u} = (|1^2 2^2|, -|1^2 2^2|, |1^2 2^2|)$
 $\frac{1}{2} \begin{matrix} 2 & 1 \\ 1 & 2 \end{matrix} = (6, -2, -2)$

$\rightarrow = \left\| \frac{(4, 1, 3) \cdot (6, -2, -2)}{(6, -2, -2) \cdot (6, -2, -2)} (6, -2, -2) \right\|$

$= \left\| \frac{16}{44} (6, -2, -2) \right\|$

$= \left\| \frac{4}{11} (6, -2, -2) \right\| = \frac{4}{11} \|(6, -2, -2)\|$

$= \frac{4}{11} \sqrt{36+4+4}$

$= \frac{4\sqrt{44}}{11}$

$= \frac{8\sqrt{11}}{11}$

Volume = base \times height

$= 6\pi \frac{8\sqrt{11}}{11} = \frac{48\pi\sqrt{11}}{11}$

Bonus Question. (5 marks) If $\vec{u}, \vec{v} \in \mathbb{R}^n$ then $|\vec{u} \cdot \vec{v}| \leq \|\vec{u}\| \|\vec{v}\|$. Hint: Analyse the squared norm of $\|\vec{u}\|\vec{v} - \|\vec{v}\|\vec{u}$ and $\|\vec{u}\|\vec{v} + \|\vec{v}\|\vec{u}$.

³From the assigned homework.