

Question 1. (1 mark each) Integrate the following indefinite integrals:

a.

$$\int \frac{1}{x^{1/3}} dx = \int x^{-1/3} dx = \frac{x^{2/3}}{2/3} + C = \frac{3x^{2/3}}{2} + C$$

b.

$$\int \sec x dx = \ln |\sec x + \tan x| + C$$


c.

$$\int \cot x dx = \ln |\sin x| + C$$

d.

$$\int \frac{1}{x\sqrt{5x^2-11}} dx = \int \frac{1}{x\sqrt{5(x^2-\frac{11}{5})}} dx$$

$$\begin{aligned} &= \frac{1}{\sqrt{5}} \int \frac{1}{x\sqrt{x^2 - (\frac{\sqrt{11}}{\sqrt{5}})^2}} dx \\ &= \frac{1}{\sqrt{5}} \frac{\sqrt{5}}{\sqrt{11}} \operatorname{arccsc}\left(\frac{x}{\frac{\sqrt{11}}{\sqrt{5}}}\right) + C \\ &= \frac{1}{\sqrt{11}} \operatorname{arccsc}\left(\frac{\sqrt{5}x}{\sqrt{11}}\right) + C \end{aligned}$$

Question 2. (5 marks) The graph of $y = f(x) = x + |\cos x|$ is given below:Find an approximation of the area under $f(x)$ on the interval $[2, 5.2]$, using the right endpoint and 4 approximating rectangles. Use the function to find the approximation and not the graph.

$$n=4$$

$$\Delta x = \frac{b-a}{n} = \frac{5.2-2}{4} = \frac{3.2}{4} = 0.8$$

$$x_i = a + i\Delta x = 2 + i(0.8)$$

$$x_1 = 2 + 1(0.8) = 2.8$$

$$x_2 = 2 + 2(0.8) = 3.6$$

$$x_3 = 2 + 3(0.8) = 4.4$$

$$x_4 = 2 + 4(0.8) = 5.2$$

$$\text{Area} \approx \sum_{i=1}^n \text{area of } R_i$$

$$= \sum_{i=1}^n f(x_i) \Delta x$$

$$= \Delta x \sum_{i=1}^n f(x_i)$$

$$= \Delta x \sum_{i=1}^n (x_i + |\cos(x_i)|)$$

$$\begin{aligned} &= 0.8 [2.8 + |\cos(2.8)| \\ &\quad + 3.6 + |\cos(3.6)| \\ &\quad + 4.4 + |\cos(4.4)| \\ &\quad + 5.2 + |\cos(5.2)|] \\ &\approx 14.89 \end{aligned}$$

Question 3. (3 marks) Determine whether the function $y = f(x)$ is a solution to the initial value problem (IVP) below. Show your work.

$$f(x) = \frac{x}{2} \arctan(2 \ln x) - \frac{\pi}{4} x.$$

Determine whether $f(x)$ satisfies

$$y' = \arctan(2 \ln x), \quad y(\sqrt{e}) = 0,$$

for $x > 0$.

$$f'(x) = \frac{\arctan(2 \ln x)}{2} + \frac{x}{2} \frac{1}{1 + (2 \ln x)^2} \cdot \frac{2}{x} - \frac{\pi}{4}$$

$$\neq y'$$

∴ does not satisfy the differential equation.
∴ is not a solution.

$$\begin{aligned} y(\sqrt{e}) &= f(\sqrt{e}) = \frac{\sqrt{e}}{2} \arctan(2 \ln \sqrt{e}) - \frac{\pi}{4} \sqrt{e} \\ &= \frac{\sqrt{e}}{2} \arctan\left(\frac{2}{2} \ln(e)\right) - \frac{\sqrt{e} \pi}{4} \\ &= \frac{\sqrt{e}}{2} \arctan(1) - \sqrt{e} \frac{\pi}{4} \\ &= \frac{\sqrt{e}}{8} \pi - \frac{\sqrt{e}}{4} \pi \neq 0 \end{aligned}$$

∴ does not satisfy the initial condition.

Question 4. (4 marks) Sketch a region whose area is equal to $\lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{1 - \left(\frac{i}{n}\right)^2} \frac{1}{n}$ and find the exact value of the area.


$$\Delta x = \frac{1}{n} = \frac{b-a}{n} \Rightarrow 1 = b-a$$

$$x_i = \frac{i}{n} = a + i \Delta x$$

$$\frac{i}{n} = a + \frac{i}{n} \Rightarrow a = 0$$

$$\Rightarrow b = 1$$

$$f(x) = \sqrt{1 - x^2}$$



$$\text{Area} = \frac{\pi(1)^2}{4} = \frac{\pi}{4}$$